

Prescribing Information

Dovato ▼ dolutegravir 50mg/lamivudine 300mg tablets

See Summary of Product Characteristics (SmPC) before prescribing

Indication: HIV-1 in adults & adolescents above 12 years of age weighing ≥ 40 kg, with no known or suspected resistance to the integrase inhibitor class, or lamivudine.

Dosing: One tablet once daily with or without food. Use an additional 50mg tablet of dolutegravir approximately 12 hours after the dose of Dovato when co-administered with efavirenz, nevirapine, tipranavir/ritonavir, etravirine (without boosted PI), carbamazepine, oxcarbazepine, phenytoin, phenobarbital, St John's Wort or rifampicin.

Elderly: Limited data in 65+ yrs. Not recommended in patients with creatinine clearance < 50 mL/min. Caution in severe hepatic impairment. **Contraindications:** Hypersensitivity to any ingredient.

Warnings/precautions: Risk of hypersensitivity reactions. Discontinue dolutegravir and other suspect agents immediately. Risks of osteonecrosis, immune reactivation syndrome. Monitor LFTs in Hepatitis B/C co-infection and ensure effective Hepatitis B therapy. Caution with metformin: monitor renal function and consider metformin dose adjustment. Use with etravirine requires boosted PI or increased dose of dolutegravir. Use with Mg/Al-containing antacids requires dosage separation. Use with calcium, multivitamins or iron also requires dosage separation if not taken at the same time with

food. Use with cladribine or emtricitabine not recommended. When possible, avoid chronic co-administration of sorbitol or other osmotic acting alcohols (see SmPC section 4.5). If unavoidable, consider more frequent viral load monitoring. **Pregnancy/ lactation:** The safety and efficacy has not been studied in pregnancy. Before initiating dolutegravir, women of childbearing potential (WOCBP) should undergo pregnancy testing. WOCBP who are taking dolutegravir should use effective contraception. Dolutegravir should not be used during the first trimester due to the potential risk of neural tube defects, unless there is no alternative. Dolutegravir should only be used during the second and third trimester of pregnancy when the expected benefit justifies the potential risk to the foetus. Avoid breast-feeding. **Side effects:** See SmPC for full details. Headache, GI disturbance, insomnia, abnormal dreams, depression, anxiety, dizziness, somnolence, rash, pruritus, alopecia, fatigue, arthralgia, myalgia, hypersensitivity, suicidal ideation or suicide attempt, hepatitis, blood dyscrasias, acute hepatic failure, pancreatitis, angioedema, rhabdomyolysis, lactic acidosis, peripheral neuropathy. Elevations of ALT, AST and CPK. **Basic NHS costs:** £656.26 for 30 tablets (EU/1/19/1370/001). MA holder: ViiV Healthcare BV, Huis ter Heideweg 62, 3705 LZ Zeist, Netherlands. Further information available from Customer Contact Centre, GlaxoSmithKline UK Ltd, Stockley Park West, Uxbridge, Middlesex UB11 1BT.

POM S1A

Trade marks are owned by or licensed to the ViiV Healthcare group of companies.

Date of approval: July 2019

PI-2451

Adverse events should be reported. For the UK, reporting forms and information can be found at www.mhra.gov.uk/yellowcard or search for **MHRA Yellowcard** in the **Google Play** or **Apple App store**. Adverse events should also be reported to GlaxoSmithKline on 0800 221441.

Adverse events should be reported. For Ireland, adverse events should be reported directly to the HPRA; Freepost, Pharmacovigilance Section, Health Products Regulatory Authority, Earlsfort Terrace, Dublin 2, Tel: +353 1 676 4971, medsafety@hpra.ie. Adverse events should also be reported to GlaxoSmithKline on 1800 244 255.