

## Prescribing Information

### Juluca<sup>▼</sup> dolutegravir 50mg/rilpivirine 25mg tablets

See Summary of Product Characteristics (SmPC) before prescribing

**Indication:** HIV-1 in virologically suppressed adults (HIV-1 RNA <50 copies/mL) on stable ART for at least 6 months with no history of virological failure and no known resistance to any NNRTI or INI. **Dosing:** *Adults (over 18 years):* one tablet once daily **with food**.

*Elderly:* Limited data in 65+ yrs. Caution in severe hepatic or renal impairment.

**Contraindications:** Hypersensitivity to any ingredient. Co-administration with carbamazepine, oxcarbazepine, phenobarbital, phenytoin, rifampicin, rifapentine, proton pump inhibitors, systemic dexamethasone (excluding single dose), St John's Wort or fampridine.

**Special warnings/precautions:** Risk of hypersensitivity reactions. Discontinue Juluca immediately if suspected. Risks of prolongation of QTc interval, osteonecrosis, opportunistic infections, immune reactivation syndrome. Monitor LFTs in Hepatitis B/C co-infection and ensure effective Hepatitis B therapy. Small rise in serum creatinine in first 4 weeks of treatment, not considered clinically relevant. Do not co-administer with other antiretrovirals (except in case of co-administration of rifabutin, when an extra dose of rilpivirine 25mg should be used). Use with antacids or once-daily H<sub>2</sub>-receptor antagonists requires dosage separation. Calcium, iron or multivitamins should be taken at the same time as Juluca with food, otherwise dosage separation recommended. Caution with

metformin: monitor renal function and consider metformin dose adjustment to minimise risk of lactic acidosis. If macrolide antibiotics are required, consider azithromycin.

Caution with antimalarials

(artemether/lumefantrine) or anticoagulants

(dabigatran). **Pregnancy/ lactation:** Not recommended during pregnancy. Before initiating Juluca, women of childbearing potential (WOCBP) should undergo pregnancy testing. WOCBP who are taking Juluca should use effective contraception. Do not breast-feed. **Side effects:** See SmPC for full details.

Increased total and LDL cholesterol, insomnia, headache, dizziness, nausea, diarrhoea, increased triglycerides, decreased appetite, abnormal dreams, depression, anxiety, sleep disorders, GI disorders, rash, pruritus, fatigue, decreased white blood cell count, haemoglobin and platelet count, arthralgia, myalgia, hypersensitivity, hepatitis, suicidal ideation or suicide attempt, acute hepatic failure. Changes in laboratory biochemistries: elevations of ALT, AST, pancreatic amylase, bilirubin and CPK.

**Basic NHS costs:** £699.02 for 30 tablets. **MA**

**number:** EU/1/18/1282/001. **MA holder:**

ViiV Healthcare BV, Van Asch van Wijckstraat 55H, 3811 LP Amersfoort, Netherlands.

Further information available from

[customercontactuk@gsk.com](mailto:customercontactuk@gsk.com) Freephone 0800 221 441.

POM

Trade marks are owned by or licensed to the ViiV Healthcare group of companies.

Date of approval: May 2020

PI-0396v3

Adverse events should be reported. For the UK, reporting forms and information can be found at [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for **MHRA Yellowcard** in the **Google Play** or **Apple App store**. Adverse events should also be reported to GlaxoSmithKline on 0800 221441.